

Sward enhancement: diversifying grassland by spreading species-rich green hay

Sward enhancement refers to management techniques which aim to increase the botanical diversity (mainly the wildflower component) of species-poor grassland. Such work can be funded under Environmental Stewardship, in particular Higher Level Stewardship. Spreading species rich green hay is one method of sward enhancement. Other techniques of oversowing, slot seeding and planting pot-grown transplants and plugs are described in separate notes.

Key points

- Both donor and receptor sites for green hay must be carefully selected.
- The receptor site should be prepared prior to spreading to achieve a short sward with 40-50% bare ground.
- Use green hay spread within a few hours of collection.
- Hay must be bedded in after spreading (eg by trampling with stock).
- Subsequent site management is required.

Introduction

Not all grassland is suitable for enhancement. The main requirements include low soil fertility and low/no weed burden. Enhancement methods usually involve disturbance to the sward, therefore some sites may not be suitable eg where it could damage historical or bird interest, or increase the risk of soil erosion. For more information see Technical Information Note TIN061 - *Sward enhancement: selection of suitable sites* and TIN062 - *Sward enhancement: choice of methods*.

The term green hay refers to herbage cut at or just before the hay stage, which is collected without prior wilting or turning and spread immediately on the receptor site.

If collected from species rich grassland, and used correctly on the right sites, it can be a very effective method of sward enhancement.



Mown hay meadow in Lower Derwent Valley NNR

The main advantages of using green hay are that it is usually cheaper than purchasing commercial seed, and it is a good means of ensuring that fresh seed, from a local source, is used. In addition, a wider range of species may be contained in green hay than is available as seed.

Sward enhancement: diversifying grassland by spreading species-rich green hay

Green hay, rather than conventional dry hay, is recommended because a much higher proportion of the seeds remains in the flower heads. Research has found that a broad range of species is present in green hay, including early flowering species such as cowslip *Primula veris*. There are also indications that using green hay is a good method of promoting colonisation by orchids.

In dry hay, many seeds are shed in the processes of drying and collecting, and what remains is mostly grass seed rather than wildflower seed. Spreading dry hay may introduce a few wildflowers (especially if the hay is carefully handled and fresh), but it is unreliable and it is not considered suitable as the main method of sward enhancement.

To be successful, the technique of spreading green hay requires very careful organisation. The receptor site must be ready to receive the hay when the donor site is cut.

Green hay cannot be stored for more than a few hours before it heats up. Any heating will threaten the viability of the seeds. Once collected the green hay must be immediately transported to and spread on the receptor site.

Research to date has focused on the use of green hay to enhance hay meadows, and the method is probably easier to undertake on meadows than pastures. However, there are also encouraging results from experiments on the use of green hay to diversify pastures.

Terminology

Green hay	Herbage cut at or just before the hay stage, which is collected immediately without prior wilting or turning.
Donor site	Grassland used as the source of green hay.
Receptor site	Grassland on which the green hay is to be spread.

Donor site

Choice of site

Donor sites for green hay should be grassland with the following characteristics:

- It should be in the same locality as the receptor site and support the target vegetation community that is to be created on the receptor site.
- Site characteristics should be similar to the receptor site, eg soil type, soil pH, hydrology and management (meadow/pasture).
- It should be species rich.
- It must be free from pernicious weeds such as spear thistle *Cirsium vulgare*, ragwort *Senecio jacobaea* and broad-leaved dock *Rumex obtusifolius*.
- Highly competitive species such as Yorkshire fog *Holcus lanatus*, white clover *Trifolium repens* and creeping buttercup *Ranunculus repens* should not form a major part of the sward.
- It must be physically possible to cut and collect the vegetation. Steep banks may not be suitable.
- It must be large enough to supply sufficient green hay for use on the receptor site. **As a guide, material cut from 1 ha should be sufficient to spread on to 3 ha.**

Possible donor sites include species rich meadows, banks, road verges and churchyards. It is important to avoid damage to nesting birds, invertebrates and plants of interest on donor sites, particularly on sites that are not normally managed by cutting.

For advice on what is an acceptable donor site for a particular hay spreading operation, contact your local Natural England Adviser. Permission to take green hay will be required from Natural England if the site is a SSSI or is in an agri-environment scheme agreement

Frequency of use

Many grassland species typically have very poor soil seed banks, as they rely on rapid germination in the autumn or spring following seed shedding. Green hay removes more seed

Sward enhancement: diversifying grassland by spreading species-rich green hay

than dry hay and risks reducing the diversity of the donor site. Donor sites should therefore not be used too frequently. As a guide, green hay should be taken from no more than one third to one fifth of a site in any one year, and rotate cut areas each year.

Green hay collection

The sward must be left uncut and ungrazed for between 8 to 12 weeks prior to collecting green hay. This will allow the plants to flower and set seed.

Where the donor site is a traditionally managed, species rich hay meadow it should be cut at the usual time for that site (usually mid July to early August). Most plants in those meadows will have completed flowering and seeding by hay cutting time, as a consequence of years of such management.

In pastures, plants flower and set seed over a longer period and it is more difficult to collect the full range of species with a single cut. A later cutting date (eg late August/early September), which allows the seed of late flowering species such as harebell (*Campanula rotundifolia*) to be collected, may be better.

Cut material should be collected within 24 hours of cutting and should not be wilted or turned.

Receptor site

The success of hay spreading depends on the presence of gaps in the sward which are large enough for seeds to germinate and establish free from competition. Seeds must land and then be pressed into the bare soil in the gaps, with sufficient moisture to germinate and sustain them.

Weed control

All pernicious weeds present in the sward should be controlled prior to ground preparation. Any application of herbicide should be by spot treatment or weed wiping to avoid damaging non target species.

Spear thistle, ragwort and other weeds with wind-blown seeds should be controlled where they occur on adjacent areas, as they can

quickly invade once the sward has been opened up.

Ground preparation

In the majority of cases, ground preparation will be required prior to hay spreading. Spreading hay onto a closed sward is extremely unlikely to be successful. Seed may fail to come into contact with the soil and die, and any seedlings which germinate may be out competed by the existing sward.

The starting point is a short sward, created by cutting (with cuttings removed) or grazing. The aim is then to create 40-50% bare ground, using livestock (the 'hoof and tooth' method) or machinery. This may seem drastic but research and experience has shown this to be necessary to reduce competition from the existing sward. Within a few months the sward will recover and very little bare ground will remain.

Where livestock are used to create bare ground, cattle are most effective as their hooves more easily break up the sward. Sheep can also be effective on damp ground. As a guide there should be frequent gaps of at least 10 cm in diameter.

Where livestock are unavailable, or where the ground is too dry for them to be effective, bare ground can be created using a power harrow or set of discs.

Mechanical sward disturbance may release more nitrogen from the soil and stimulate the weed seed bank, so it should only be undertaken on sites with very low fertility and no weed burden. In a few cases, notably in upland hay meadows, there may be sufficient bare ground and soil moisture following the hay cut for seed to be sown without further ground preparation. However, such cases are likely to be rare.

Spreading the hay

Once the cut material has been collected from the donor site, it must be transported to and spread on the receptor site on the same day - ideally within an hour or two. It is essential that it does not heat up.

Sward enhancement: diversifying grassland by spreading species-rich green hay

The hay should be spread thinly and evenly so that it does not create a mulch which will inhibit seed germination. The sward should be visible beneath the hay.

Leave the hay for at least one week in dry weather, or three weeks in wet weather, to allow seed to fall. After this period allow stock (ideally cattle) to graze the site, and trample out and bed in the seeds. Alternatively, roll the site lightly.

Remove the hay if it is smothering the sward. However, this is unlikely to occur if hay is spread in the recommended amounts.

Successive hay spreading

If hay spreading has been undertaken correctly using a species rich donor site, it should not be necessary to repeat the operation. Major disturbance to the sward within at least the first 2 years should be avoided, to allow plants to establish. Some of the introduced species may not appear in the sward for several years, so the success of hay spreading should not be judged too soon.

Methods of collecting and spreading green hay which have been used successfully include:

Large areas

Use a direct cut forage harvester, discharging into a trailer or muck spreader. Spread the herbage with the muck spreader, or fork it out of the trailer.

Cut with a mower. (A mower conditioner should not be used as this will encourage seed drop at the donor site.) Then either:

- Fork the herbage into a trailer and fork it out onto the receptor site.
- Gently row up the herbage, then bale it with a round baler into loose bales. Bales should be tied or netted but not wrapped. (Note that green bales will be considerably heavier than bales of dry hay. To make spreading easier, the thickness of the layers forming the bales can be reduced by decreasing the forward speed of the baler whilst maintaining PTO speed.) Spread the bales by:

- Unrolling and forking them out by hand.
- Unrolling them, then spreading them with a hay bob/tedder.
- Using a straw chopper.
- Using a muck spreader.
- Gently row up the herbage, then collect it with a forage harvester discharging into a muck spreader or a trailer. Spread the chopped material with the muck spreader, or fork it out of the trailer.



Spreading green hay by muck spreader

Small areas

Cut the vegetation with a brush cutter or power scythe. Then either:

- Fork the herbage into a trailer and fork it out onto the receptor site.
- Row up the herbage and bale it with a mini baler. Unroll the bales and fork them out onto the receptor site.

Subsequent management

In the period immediately after hay spreading (usually July - November), the sward should be kept short so that light can aid germination. This is best done by short periods of intensive grazing. Alternatively the sward can be cut and the cuttings removed. Prolonged grazing should initially be avoided in order to reduce the risk of seedlings being selectively grazed.

In the first spring, it may be necessary to cut or graze the sward to avoid seedlings being shaded out by the existing vegetation. A short period of intensive grazing, or cutting (with the

Sward enhancement: diversifying grassland by spreading species-rich green hay

cuttings removed) is recommended. However, this may not be appropriate on all sites and care should particularly be taken to avoid damage to other interests on the site eg birds and invertebrates.

Any perennial weeds which have colonised should be controlled early on, eg by spot treatment with herbicide. Any annual weeds are likely to be controlled by the regular cutting or grazing outlined above.

In subsequent years, if the field is to be managed as a hay meadow it should be cut late (eg after mid July), with swath turning or tedding undertaken to assist seed shedding. The cutting date should be matched to that of the donor meadow, if possible. The use of livestock, particularly for aftermath grazing, is important because they create gaps in the sward and trample in the seed, which helps the introduced species to spread.

Where the field is managed as pasture, plants must be allowed to flower and set seed by reducing the grazing pressure for a period of about eight weeks in the spring and summer.

There should be no use of inorganic fertilisers or widespread application of herbicides.

Commitment to the appropriate long term management is essential if the grassland enhancement is going to succeed and be maintained.

Further information

Natural England Technical Information Notes are available to download from the Natural England website: www.naturalengland.org.uk. In particular see:

- Technical Information Note TIN035: *Soil sampling for habitat recreation and restoration in agri-environment schemes*

- Technical Information Note TIN036: *Soils and agri-environment schemes: interpretation of soil analysis*
- Technical Information Note TIN038: *Seed sources for grassland restoration and re-creation in Environmental Stewardship*
- Technical Information Note TIN050: *Selecting indicators of success for grassland enhancement*
- Technical Information Note TIN060: *The use of yellow rattle to facilitate grassland diversification in agri-environment schemes;*
- Technical Information Note TIN061: *Sward enhancement: selection of suitable sites*
- Technical Information Note TIN062: *Sward enhancement: choice of methods*
- Technical Information Note TIN064: *Sward enhancement: diversifying grassland by oversowing and slot seeding*
- Technical Information Note TIN065: *Sward enhancement: diversifying grassland using pot-grown wildflowers or seedling plugs*

For further information contact the Natural England Enquiry Service on 0845 600 3078 or e-mail enquiries@naturalengland.org.uk.

This note does not supersede prescriptions in agri-environment scheme agreements. If there is any conflict between the information in this note and your agreement please contact your local adviser.

Authors and contributors

This note was previously published as RDS Technical Advice Note 28. It was compiled by Morwenna Christian and Steve Peel, with major contributions by Marion Bryant. Much of the content of this note is informed by Defra funded research by the University of Reading, and by the work of Professor Ian Trueman (University of Wolverhampton). Editor Susie Smith, photographs Peter Roworth and Alistair Helliwell.